



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶: A61K 45/00, 45/05, 38/20	A1	(11) International Publication Number: WO 96/10423 (43) International Publication Date: 11 April 1996 (11.04.96)
(21) International Application Number: PCT/US95/12611 (22) International Filing Date: 28 September 1995 (28.09.95) (30) Priority Data: 08/316,622 30 September 1994 (30.09.94) US 08/408,915 22 March 1995 (22.03.95) US (71) Applicants: LUDWIG INSTITUTE FOR CANCER RE- SEARCH [CH/US]; 1345 Avenue of the Americas, New York, NY 10105 (US). SLOAN-KETTERING INSTITUTE FOR CANCER RESEARCH [US/US]; 1275 York Avenue, New York, NY 10021 (US). (72) Inventors: NOGUCHI, Yuji; 1275 York Avenue, New York, NY 10021 (US). CHEN, Yao-tseng; Cornell Medical Center, York Avenue, New York, NY 10021 (US). OLD, Lloyd, J.; 1345 Avenue of the Americas, New York, NY 10105 (US). (74) Agent: HANSON, Norman, D.; Felfe & Lynch, 805 Third Avenue, New York, NY 10022 (US).		(81) Designated States: AU, CA, CN, FI, JP, NO, NZ, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the</i> <i>claims and to be republished in the event of the receipt of</i> <i>amendments.</i>
(54) Title: COMPOSITIONS CONTAINING A p53 DERIVED PROTEIN OR PEPTIDE, AN ADJUVANT, AND INTERLEUKIN-12 AND USES THEREOF		
(57) Abstract The invention involves the combination of Interleukin-12 with p53 derived peptides and an adjuvant, preferably QS-21. It is found that this combination provokes a surprisingly strong immune response. Further, in an accepted <i>in vivo</i> model, the use of compositions containing these three ingredients led to diamatic decreases in the growth of induced tumors, thus suggesting a therapeutic regime.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

5 COMPOSITIONS CONTAINING A p53 DERIVED PROTEIN
 OR PEPTIDE, AN ADJUVANT, AND INTERLEUKIN-12
 AND USES THEREOF

RELATED APPLICATION

10 This application is a continuation-in-part of patent
application Serial No. 08/316,622, filed on September 30,
1994.

FIELD OF THE INVENTION

15 This invention relates to compositions useful in the
generation of an immune response. The immune response
includes, inter alia, provocation of T cells, such as
cytolytic T cells, against complexes of a p53 derived peptide
and MHC molecules or a p53 derived peptide or protein itself,
as well as antibodies against such peptides and proteins.
Such T cells and antibodies may be generated, e.g., in a
20 mouse, rat, rabbit, sheep, goat or other non-human animal, and
then used in diagnostic methods to identify tumor presence.
They may also be generated in vitro such as by cell culture
and hybridoma techniques. The compositions may also be used,
therapeutically, via administration to a subject afflicted
25 with a cancerous condition or one where cell transformation
has taken place, to provoke an immune response against tumors,
cancer cells, and transformed cells.

BACKGROUND AND PRIOR ART

30 Adjuvants, broadly defined, are substances which promote
immune responses. Frequently, the adjuvant of choice is
Freund's complete adjuvant, or killed B. pertussis organisms,
used in combination with alum precipitated antigen. A general
discussion of adjuvants is provided in Goding, Monoclonal
Antibodies: Principles & Practice (Second edition, 1986), at
35 pages 61-63, which are incorporated by reference herein.
Goding notes, however, that when the antigen of interest is of
low molecular weight, or is poorly immunogenic, coupling to an
immunogenic carrier is recommended. Such molecules, according
to Goding, generally have molecular weights below about 1000.
40 Among the carriers suggested by Goding, at page 283, are

5 keyhole limpet hemocyanin, bovine serum albumin, ovalbumin, and fowl immunoglobulin.

What is problematic about such carriers, however, is that frequently they are also immunogenic themselves. Thus, the immune response may be a general one, with part, most, or all of it being directed against the carrier molecule rather than the immunogen itself.

Exemplary of developments in the art as they relate to adjuvants is U.S. Patent No. 5,057,540 to Kensil, et al, the disclosure of which is incorporated by reference herein. Kensil et al disclose the preparation of various saponin extracts, which are useful as adjuvants in immunogenic compositions. As they are natural products, the extracts are not completely defined. Kensil, et al do provide a complete and enabling disclosure for how various extracts, including QA-7, QA-19, and QA-21 (also referred to as QS-21) are prepared. Experiments are set forth in which bovine serum albumin ("BSA") was combined with various extracts (examples 8 and 9), and where feline leukemia virus recombinant glycoprotein "gp70RΔ" was tested, following absorption to aluminum hydroxide (alum). The two immunogens tested, however, are expected to be immunogenic in their own right (gp70RΔ has a molecular weight of 70 kd, and serum albumin has about the same molecular weight). No experiments were carried out at all on molecules which should, per se, be considered to be poorly or even non-immunogenic, and thus would be expected to require the use of alum absorption or the use of haptenic carriers for provocation of a response.

In PCT Application WO9219758, which corresponds to defensive publication 7697275, which is incorporated by reference herein, an adjuvant referred to as "MTP-MF59" is disclosed. This adjuvant is used in connection with a Plasmodium falciparum protein, "Pfs-25-B". This combination is described as a transmission blocking vaccine. The P. falciparum protein is itself large enough to be immunogenic. Thus, none of the art shows that the improved adjuvants can be used in combination with presumptively non-immunogenic

5 proteins and peptides to yield immunologically effective compositions.

Interleukin-12, or "IL-12" hereafter, is known to play an important role in the differentiation of T_H0 - T_H1 , and T_H2 - T_H0 cells in vivo. See Manetti, et al, J. Exp. Med 177:1199-1204
10 (1993); Hsieh, et al, Science 260:547-549 (1993).

The molecule referred to as p53 has been known for a number of years. A brief summary of the molecule is provided by Culotta, et al., Science 262: 1958-1961 (1993) ("p53 Sweeps Through Cancer Research"). As reported therein, about 50% of
15 all subjects diagnosed with cancer have a p53 mutation (or p53 mutations). These mutations can abolish the function of normal, or wild type p53, which is to act as a tumor suppressor of the observed mutations, about 90% involve a change in amino acid sequence (a "missense" mutation), as
20 compared to "nonsense" mutations, where the changes in the gene result in truncation or destabilization of the protein. The wt p53 molecule is also involved in the cell cycle, in that it inhibits cell division. This is accomplished by activating a second protein of 21 kilodaltons, which inhibit
25 the proteins "cdk" and "cyk", which are required for cell division.

It has now been found, surprisingly, that compositions can be made which comprise immunogenic peptides, such as p53 derived peptides, in combination with an adjuvant such as
30 those described above and the cytokine interleukin-12, or "IL-12". When administered to a subject animal, these compositions provoke an immunogenic response which is surprisingly strong, and totally unexpected in view of the response elicited with each element by itself, or pairs of
35 these elements. In especially preferred embodiments the immunogenic peptide or protein is combined with the adjuvant QS21, which is disclosed in the Kensil, et al, patent, incorporated by reference supra.

A further aspect of this invention is the use of the
40 cytokine, IL-12. Other interleukins, i.e., IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, and IL-11, may also

5 be used in view of their known efficacy as T-cell and B-cell growth factors. Of these, IL-2 is especially preferred. The compositions yield unexpected results when administered to T cells or B cells, in vitro or in vivo, in that their proliferation or other response, is surprisingly enhanced.

10 The immunogens of this invention include all p53 derived peptides and proteins, such as mutant p53 peptides, as exemplified.

The invention is described in greater detail in the disclosure which follows.

15 **BRIEF DESCRIPTION OF THE FIGURES.**

Figures 1A-1E, inclusive, depict experiments showing the antitumor effect of IL-12. In these figures, differing doses of IL-12 were administered to mice which had been injected with Meth A sarcoma cells.

20 Figures 2A-2C show the importance of the timing of the administration of IL-12. In figure 2A, IL-12 was administered 14 days before challenge with Meth A cells. In Figure 2B, the IL-12 was administered on the same day that the challenge took place. In Figure 2C, the IL-12 was administered 7 days after challenge.

25 Figures 3A-3J cover the use of two adjuvants, QS-21 and Incomplete Freund's adjuvant, together with IL-12. The experiments are directed to the study of IL-12's immunomodulatory effect. In Figures 3A-3E, QS-21 was used. In Figures 3F-3J, IFA was used. As explained infra, various doses of IL-12 were used.

30 Figures 4A-4C show that there was strong suppression of tumor growth when low doses of IL-12 were administered with peptides. In Figure 4A, the mutant peptide was administered in QS-21. In Figure 4B, wild type peptide was combined with QS-21, and in Figure 4C, QS-21 was used alone. The peptide/adjuvant combination was administered 7 and 14 days after challenge with peptides. A very low dose of IL-12 (1 ng) was administered at day 7, and then as elaborated upon infra.

40

5 **DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS****Example 1**

The anti-tumor effect of IL-12, used in combination with mutant p53 peptides, was studied.

10 The in vivo model employed used the transplantable, methylcholanthrene-induced sarcoma "Meth A." This is a sarcoma of BALB/C (murine) origin, passaged as an ascitic tumor. See Old, et al, Ann. N.Y. Acad. Sci 101:80-106 (1962), incorporated by reference in its entirety.

15 BALB/C female mice were injected intradermally with 5×10^4 Meth A cells on day 0. Seven days later, a treatment regime began, wherein the animals received doses of IL-12, 4 times a week. Control animals received no IL-12. The remaining animals received 1 ng, 10 ng, 100 ng, or 500 ng of IL-12 in each dose, administered by intraperitoneal injection.

20 The effect of the IL-12 on the induced tumors was observed by measuring the diameter of the tumors at the end of the treatment period. Figures 1A, 1B, 1C, 1D and 1E show this. The panels show results with doses of 0 ng, 1 ng, 10 ng, 100 ng, and 500 ng, respectively.

25 It is clear from these figures that there is a dose dependent anti-tumor effect provoked by IL-12, although the doses are extremely small.

30 The effect of the IL-12 was abolished by administering monoclonal antibodies against CD8 molecules, but was not affected via administration of anti-CD4 mAbs. This suggests that cytolytic T cells are an essential part of the response.

Example 2

35 The importance of timing in the administration of IL-12 was then studied. In these experiments, female BALB/C mice were again injected with 5×10^4 Meth A cells, as in Example 1 (day 0). Injections of 100 ng IL-12 were administered at day "-14" (i.e., 14 days before challenge with the tumor cells), day 0, or day 7 (i.e., 7 days after challenge) as in Example 1. These results, presented in figures 2A (day "-14"), 2B (day 0), and 2C (day 7) show that IL-12 is most
40 effective after the tumor had become established.

5 Example 3

The experiments set forth in this example evaluate the effect of IL-12 on generation of cytolytic T cells ("CTLs" hereafter) specific to peptide/MLTC complexes.

10 Peptide 234 CM has amino acid sequence:

Lys Tyr Ile Cys Asn Ser Ser Cys Met (SEQ ID NO: 1)
and contains a point mutation in codon 234 of Meth A p53. The wild type counterpart is

Lys Tyr Met Cys Asn Ser Ser Cys Met (SEQ ID NO: 2)
15 BALB/C female mice were immunized with 100 µg of the wild type peptide, combined with the adjuvant QS-21, described supra, or Incomplete Freund's Adjuvant ("IFA"). A total of 10 µg of QS-21 was used for each injection. The IFA was dissolved in PBS, at concentrations of 1 mg/ml. The combination of peptide and
20 adjuvant was administered twice at weekly intervals. Over the two week period, the animals were also injected with IL-12, four times. The dose of IL-12 was either 0.1 ng, 1 ng, 10 ng, or 100 ng. Control animals received no IL-12.

One week after the last injections, the spleen cells of
25 the immunized mice were sensitized with syngeneic spleen cells, pulsed with the mutant peptide (234CM). Sensitization was carried out using well known techniques. The activity of CTLs in the sample was determined using the well known ⁵¹Cr release assay. This methodology is described
30 by Noguchi, et al, Proc. Natl. Acad. Sci USA 91:3171-3175 (1994), incorporated by reference. Sensitization with pulsed spleen cells is a methodology which leads to a much stronger response than if peptide alone were administered.

The results are presented in figure 3, where figures 3A
35 and 3F present data from controls, figures 3B and 3G show the data obtained when 0.1 ng of IL-12 was used, figures 3C and 3H show results where 1 ng of IL-12 was used, figures 3D and 3I present the results where 10 ng of the cytokine were used, and 3E and 3J, where 100 ng were used. Data are presented in
40 terms of the percentage of chromium release, where the amount of peptide contacted to the cells (2.5 µg or 10 µg), figures

5 3A-3E are from experiments in which QS-21 was used, while figures 3F-3J resulted from experiments where IFA was used.

In figures 3A-3J, the filled in dots (●-●) represent the data from the mutant peptide experiments. Also presented are results in which the wild type peptide was pulsed to cell line
10 PI-HTR (a cell line derived from the mastocytoma P815, of DBA/2 origins. (See Old, et al, supra). These data are filled in squares (■-■). Also presented are data where PI-HTR were pulsed in the absence of peptide (▲-▲). This model
15 is used, and is useful because one of the known features of Meth A is that it does not generate a CTL response. PI-HTR, however, does, and can be used as a model to determine the efficacy of the compositions discussed herein.

The data presented in figure 3 show that 1 ng of IL-12 was the optimal dose for the generation of CTLs, while higher
20 doses had a suppressive effect. No CTLs were elicited following immunization with wild type peptide, regardless of the adjuvant or treatment with IL-12.

Example 4

This set of experiments investigated whether the
25 combination of a peptide immunogen and a low dose of IL-12 would suppress the growth of established tumors.

As with the prior experiments, female BALB/C mice were injected with 5×10^4 Meth A cells at day 0. The mice received two injections of the mutant peptide in QS-21, the
30 wild type peptide in QS-21, or QS-21 alone in the same manner described in Example 3, at weekly intervals. At day 7, the mice began receiving injections of 1 ng of IL-12 (4 times a week, for two weeks).

The results are presented in figures 4A-4C. Figure 4A
35 resulted from QS-21 plus the mutant peptide. Figure 4B resulted from QS-21 plus peptide, followed by injections with IL-12. Figure 4C resulted from QS-21 alone. These figures show that the IL-12 had a growth suppressive effect on the tumors. In results not depicted, immunization with wild type peptide
40 did not lead to suppression of tumor growth, even when combined with QS-21 and IL-12.

5 Also not shown are results in which QS-21 was shown to be more effective than IFA in inducing the regression of the tumors.

10 The foregoing examples demonstrate a composition comprising an amount of a p53 derived peptide, an adjuvant, e.g., QS-21, and Interleukin-12, as well as the use of the composition in the in vivo treatment of cancer. The immunogen is used in an amount sufficient to provoke an immune response against tumor cells which present it on their surface, and combined with QS-21 and IL-12. Among the noteworthy features of the invention is the fact that none of the peptide, the adjuvant, nor the IL-12 worked to stimulate an immune response when used alone, nor was the response particularly strong when the peptide was combined with adjuvant. Further, the results show that the IL-12 is particularly effective after tumors have already formed, and as combined with the peptide and the adjuvant, is effective in very small doses.

20 The compositions of the invention comprise any peptide or protein, derived from p53, in combination with a pharmaceutically acceptable adjuvant. Preferred embodiments of the invention utilize the peptides of SEQ ID NO: 1 and 2. Other p53 based peptides may also be used.

30 As will be seen from the foregoing discussion, an important aspect of the invention is stimulation of proliferation of T cells. This can be an initial stimulation or an augmentation of a prior stimulation. In particular, it is desirable to stimulate cytolytic T cells. The cytolytic T cells recognize complexes of MHC and peptide, bind thereto via their receptor, and proliferate. They also lyse the recognized cells. This response can be used not only in vivo, but in vitro, as it is well established that cytolytic T cells specific for particular complexes of MHC and peptide are present in the blood of subjects who have experienced cell transformation. By contacting a blood sample of individuals in vitro with the peptide of interest and cells which present the MHC molecule of interest, any cytolytic T cells in the blood sample will expand, i.e., proliferate. This

5 proliferation can be measured via any of the well known assays designed therefor. Especially preferred are the well known radioactive chromium (^{51}Cr) release assay, and the measurement of release of tumor necrosis factor (TNF). These assays are efficacious when peptides such as mutant p53 peptides and
10 other p53 peptides, are used.

The compositions are also useful as stimulators of B cell proliferation, or antibody production. Again, it is well known that B cells produce antibodies, and the size of their targets are well within the sizes of the p53 antigen. As with
15 T cells, the stimulation may be "ab initio", or an augmentation of a prior response, in vitro or in vivo.

The amount of peptide used will vary, depending upon the purpose of the immunization and the subject to which it is administered. For example, in the case of generating murine
20 antibodies which can then be used, e.g., to diagnose for the presence of cancer cells presenting a p53 peptide, the amount of protein or peptide may be less than that used in a course of in vivo therapy, such as that described in the example, supra. In general, a preferred dose can range from about 1 ug
25 to about 750 ug of protein or peptide per dose. In a preferred embodiment, the range is from about 10 ug to about 500 ug. Most preferably, anywhere from about 30 ug to about 300 ug per dose may be used. Of course, in the context of the therapeutic aspect of the invention, the investigator will
30 modify the dose, as a six month old infant will require dosing different from a full grown man, e.g. The mode of administration may vary, with preferred forms being oral, subcutaneous, intramuscular, intravenous and intraperitoneal administration.

35 The choice of protein or peptide in the composition will depend upon parameters determinable by the artisan. It is art recognized, for example, that different peptides are presented by the various MHC molecules. As such, if a subject is typed, using well known techniques, as presenting HLA-A2 molecules on
40 the surface of tumor cells, one will use a peptide presented by HLA-A2 molecules rather than one presented by, e.g., HLA-

5 Cw* 1601. Similarly, using techniques such as polymerase
chain reaction ("PCR"), lysis studies, and other assay
methodologies which are well known in the art, one can
determine which protein or peptide is being expressed by a
subject patient. This will lead to the decision as to what
10 protein or peptide to use. Again, by way of example, if a
subject's tumor cells are expressing mutant p53 but not normal
p53 peptide, the peptide used in immunization should be
derived from the mutant form.

Other aspects of the invention will be clear to the
15 skilled artisan and need not be reiterated here.

The terms and expressions which have been employed are
used as terms of description and not of limitation, and there
is no intention in the use of such terms and expressions of
excluding any equivalents of the features shown and described
20 or portions thereof, it being recognized that various
modifications are possible within the scope of the invention.

5

(1) GENERAL INFORMATION:

(i) APPLICANTS: Noguchi, Yugi; Chen, Yao-tseng;
Old, Lloyd J.

10

(ii) TITLE OF INVENTION: Compositions Containing A p53
Derived Protein Or Peptide, An Adjuvant, And
Interleukin-12 And Uses Thereof

(iii) NUMBER OF SEQUENCES: 2

(iv) CORRESPONDENCE ADDRESS:

15

(A) ADDRESSEE: Felte & Lynch

(B) STREET: 805 Third Avenue

(C) CITY: New York City

(D) STATE: New York

(E) COUNTRY: USA

(F) ZIP: 10022

20

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Diskette, 5.25 inch, 360 kb
storage

(B) COMPUTER: IBM PS/2

(C) OPERATING SYSTEM: PC-DOS

25

(D) SOFTWARE: Wordperfect

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER: 08/408,915

(B) FILING DATE: 22-MARCH-1995

(C) CLASSIFICATION: 435

30

(vii) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: 08/316,622

(B) FILING DATE: 30-SEPTEMBER-1994

(viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: Hanson, Norman D.

35

(B) REGISTRATION NUMBER: 30,946

(C) REFERENCE/DOCKET NUMBER: LUD 5391.1

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: (212) 688-9200

(B) TELEFAX: (212) 838-3884

40

12

5 (2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 9 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

Lys Tyr Ile Cys Asn Ser Ser Cys Met

5

15

(2) INFORMATION FOR SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 9 amino acids

20 (B) TYPE: amino acid

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

Lys Tyr Met Cys Asn Ser Ser Cys Met

25

5

5 We claim:

1. Composition useful in provoking an immune response comprising an amount of (i) a p53 derived protein or peptide, (ii) an adjuvant, and (iii) Interleukin-12, sufficient to provoke an immune response.

10 2. The composition of matter of claim 1, wherein said protein or peptide binds to an MHC molecule.

3. The composition of claim 1, wherein said peptide or peptide is derived from a protein expressed by a transformed cell.

15 4. The composition of claim 3, wherein said protein or peptide is a mutant p53 protein or peptide.

5. The composition of claim 1, wherein said immune response is a B cell response.

20 6. The composition of claim 1, wherein said adjuvant comprises a saponin.

7. The composition of claim 6, wherein said adjuvant comprises QS-21.

8. Kit useful in provoking an immune response, comprising a container means and a separate portion of each of

25 (i) a p53 derived protein or peptide,
 (ii) an adjuvant, and
 (iii) Interleukin-12.

9. Kit useful for provoking an immune response, comprising a container means which holds a separate protein of
30 each of

 (i) a nucleic acid molecule which codes for a molecule which is processed in a transformed cell into at least one p53 derived protein or peptide,

35 (ii) an adjuvant, and
 (iii) Interleukin-12.

10. The kit of claim 8, wherein said nucleic acid molecule codes for a protein or peptide associated with cell transformation.

40 11. Method for provoking an immune response in a subject, comprising administering to said subject an immune stimulating amount of

- 5 (i) a p53 derived protein or peptide,
 (ii) an adjuvant, and
 (iii) Interleukin-12.





12. The method of claim 11, wherein said immune response
is a T cell response.

10 13. The method of claim 11, wherein said adjuvant is QS-
21.





14. The method of claim 11, wherein said protein or
peptide is associated with cell transformation.

15 15. The method of claim 11, wherein said subject is
afflicted with a tumor.





1/5

FIG. 1A**FIG. 1B****FIG. 1C****FIG. 1D**





2/5

FIG. 1E**FIG. 2A****FIG. 2B****FIG. 2C**






3/5

FIG. 3A**FIG. 3B****FIG. 3C****FIG. 3D**

4/5

FIG. 3E**FIG. 3F****FIG. 3G****FIG. 3H**

5/5

FIG. 3I**FIG. 3J****FIG. 4A****FIG. 4B****FIG. 4C**

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US95/12611

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : A61K 45/00, 45/05, 38/20

US CL : 424/277.1, 85.2, 279.1

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/277.1, 85.2, 279.1

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, Medline, Biosis, Embase, Derwent

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO, A, 94/02167 (THE TRUSTEES OF PRINCETON UNIVERSITY) 03 FEBRUARY 1994, see pages 1, 4-8, and 12.	1-4 and 6-15
Y	WO, A, 94/16716 (VIROGENETICS CORPORATION) 04 AUGUST 1994, see pages 14, 22, and 25.	1-4 and 6-15
Y	Science, Volume 263, issued 25 March 1994, S.S. Hall, "IL-12 Holds Promise Against Cancer, Glimmer of AIDS Hope", pages 1685-1686, see page 1686.	1-4 and 6-15
Y	Immunology Letters, Volume 40, issued 1994, Nijman et al, "p53, A Potential Target for Tumor-Directed T Cells", pages 171-178, see pages 175-176.	2



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	*T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X*	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y*	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*G*	document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means		
P document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search 28 DECEMBER 1995	Date of mailing of the international search report 29 JAN 1996
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230	Authorized officer JULIE KRSEK-STAPLES Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORTInt. l. application No.
PCT/US95/12611**C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	Current Opinion in Immunology, Volume 4, issued 1992, P.O. Livingston, "Construction of Cancer Vaccines with Carbohydrate and Protein (Peptide) Tumor Antigens", pages 624-629, see page 627.	6, 7, and 13
Y	Human Gene Therapy, Volume 5, No. 9, issued September 1994, Wills et al, "Development and Characterization of Recombinant Adenoviruses Encoding Human p53 for Gene Therapy of Cancer", pages 1079-1088, see pages 1080, 1085-1086.	3, 9, 10, and 14